




Commonwealth Environment Research Facilities

Marine and Tropical Sciences Research Facility



The effects of Severe Tropical Cyclone *Larry* on rainforest vegetation and understorey microclimate adjacent to powerlines, highways and streams in the Wet Tropics World Heritage Area

Catherine Pohlman and Miriam Goosem



Australian Government
Department of the Environment,
Water, Heritage and the Arts

The effects of Severe Tropical Cyclone *Larry* on rainforest vegetation and understorey microclimate adjacent to powerlines, highways and streams in the Wet Tropics World Heritage Area

Catherine L. Pohlman and Miriam Goosem

School of Earth and Environmental Sciences, James Cook University



Australian Government

**Department of the Environment,
Water, Heritage and the Arts**

Supported by the Australian Government's
Marine and Tropical Sciences Research Facility
Project 4.9.3 Impacts of urbanisation on North Queensland
environments: management and remediation

© James Cook University

ISBN 978-1-921359-12-5

This report should be cited as:

Pohlman, C. and Goosem, M. (2007) *The effects of Severe Tropical Cyclone Larry on rainforest vegetation and understorey microclimate adjacent to powerlines, highways and streams in the Wet Tropics World Heritage Area*. Report to the Marine and Tropical Sciences Research Facility. Reef and Rainforest Research Centre Limited, Cairns (47pp.).

Made available online by the Reef and Rainforest Research Centre on behalf of the Australian Government's Marine and Tropical Sciences Research Facility.

The Australian Government's Marine and Tropical Sciences Research Facility (MTSRF) supports world-class, public good research. The MTSRF is a major initiative of the Australian Government, designed to ensure that Australia's environmental challenges are addressed in an innovative, collaborative and sustainable way. The MTSRF investment is managed by the Department of the Environment, Water, Heritage and the Arts (DEWHA), and is supplemented by substantial cash and in-kind investments from research providers and interested third parties. The Reef and Rainforest Research Centre Limited (RRRC) is contracted by DEWHA to provide program management and communications services for the MTSRF.

This publication is copyright. The Copyright Act 1968 permits fair dealing for study, research, information or educational purposes subject to inclusion of a sufficient acknowledgement of the source.

The views and opinions expressed in this publication are those of the authors and do not necessarily reflect those of the Australian Government or the Minister for the Environment, Water, Heritage and The Arts.

While reasonable effort has been made to ensure that the contents of this publication are factually correct, the Commonwealth does not accept responsibility for the accuracy or completeness of the contents, and shall not be liable for any loss or damage that may be occasioned directly or indirectly through the use of, or reliance on, the contents of this publication.

This report is available for download from the Reef and Rainforest Research Centre Limited website:
http://www.rrrc.org.au/publications/research_reports.html



March 2007

Contents

Executive Summary	ii
List of Figures.....	iv
List of Tables.....	iv
Abbreviations Used In This Report.....	iv
Glossary	v
Acknowledgements	vi
Introduction	1
Natural and anthropogenic disturbance near linear canopy openings.....	1
Severe Tropical Cyclone <i>Larry</i>	1
Research Questions	2
Methodology.....	3
Study Sites.....	3
Vegetation Damage.....	6
Understorey Microclimate	7
<i>Traverse Measurements</i>	7
<i>Diurnal Variation</i>	9
Results	12
Vegetation Damage.....	12
<i>Tree Damage</i>	12
<i>Sapling Damage</i>	15
Understorey Microclimate	17
<i>Traverse Measurements</i>	17
<i>Diurnal Variation</i>	19
Discussion	25
Vegetation Damage.....	25
<i>Effect of proximity to linear clearing</i>	25
<i>Effect of tree size and successional status</i>	25
<i>Effects of ‘sealed’ edges</i>	26
Understorey Microclimate	27
Cyclone <i>Larry</i> and Linear Canopy Openings.....	28
Recommendations	30
References	31
Appendix 1: Wood densities of tree species encountered in the vegetation survey ...	36

Executive Summary

Project Objectives

Severe Tropical Cyclone *Larry* crossed the coast of Queensland on 20 March 2006, damaging a large swathe of rainforest to the west of Innisfail in the Wet Tropics World Heritage Area. Within the path of the most destructive core of Cyclone *Larry* were sites previously established for a study on the effects of human-made (powerlines and highways) and natural (streams) linear canopy openings on adjacent rainforest plant communities.

Measurements of vegetation damage and understorey microclimate (light intensity, air temperature and humidity, wind speed, soil temperature and soil moisture) were made six months following the passage of Cyclone *Larry* and were compared with measurements made at the same study sites prior to the cyclone. The purposes of these measurements were to:

- Determine whether cyclone damage to vegetation was greater near the edges of human-made and natural linear canopy openings than in the forest interior; and
- Examine the influence of this vegetation damage (and consequently changes in rainforest canopy structure) on the microclimatic regimes experienced in the rainforest understorey.

Key Findings

1. Vegetation damage was slightly reduced near powerline edges, slightly elevated at intermediate distances from highway edges, and slightly elevated near stream edges, relative to the forest interior.
2. Changes in the understorey microclimatic regime mirrored the degree of damage to vegetation.
3. Where vegetation damage was greater, the understorey microclimate was brighter, warmer, drier and windier than below less-damaged areas of the forest canopy.
4. It is possible that the elevated damage at stream edges was due to a combination of two factors. Firstly, flood disturbance was noticeable: a number of trees near the edge were washed out by flood waters. Secondly, the species composition and forest structure near stream edges may be more vulnerable to cyclone damage, as stream edges have a greater proportion of pioneer species. Pioneer species suffered greater damage than mid- or late-successional tree species and these species also generally have a lower canopy and subcanopy foliage density than those commonly found at forest edges near powerlines or highways.
5. Prior to the cyclone, forest edges near powerlines and highways had developed a greater degree of edge 'sealing' than forest near stream edges (i.e. greater canopy and subcanopy foliage density) and thus reduced light availability and reduced recruitment of pioneers, weeds and other light-loving species. These 'sealed' edges may have been less prone to cyclone damage or may have offered some protection to the forest near the edge.

Management Implications and Recommendations

1. The 'sealed' forest edges near powerlines and highways appear to have offered the forest edge some degree of protection from the more severe wind damage observed near creek edges. Therefore we recommend that, for infrastructure corridors where it is not possible to minimise the footprint of the clearing (for example by raising the

powerline above the canopy), such 'sealed' edge development should be encouraged. This could be achieved through minimising disturbance to the forest edge during infrastructure and easement maintenance.

2. Unfortunately, the microclimatic changes following Cyclone *Larry* have favoured the germination and growth of numerous pioneer and weed species. Initial observations suggest that the abundance of exotic plant species within the rainforest understorey has increased substantially since the cyclone. It is recommended that this situation should be monitored to identify potential problems.

List of Figures

Figure 1:	Study areas in Wooroonooran National Park and South Johnstone State Forest.....	4
Figure 2:	Examples of linear canopy openings.....	5
Figure 3:	Instruments used for traverse measurements.....	9
Figure 4:	Instruments used for measurements of diurnal variation.....	11
Figure 5:	Covariates of cyclone damage to trees	13
Figure 6:	Cyclone damage to trees.....	14
Figure 7:	Covariate of cyclone damage to saplings.....	15
Figure 8:	Cyclone damage to saplings	16
Figure 9:	Photosynthetically active radiation	17
Figure 10:	Relative soil temperature.....	18
Figure 11:	Proportional soil moisture.....	19
Figure 12:	Diurnal temperature range and diurnal VPD range	21
Figure 13:	Maximum wind speed.....	22
Figure 14:	Edge gradients in air temperature and VPD in the morning, at midday and in the afternoon.....	24

List of Tables

Table 1:	Damage categories for trees and saplings.....	7
Table 2:	Cyclone damage to trees.....	12
Table 3:	Cyclone damage to saplings	15
Table 4:	Tests of between-subject effects for MANOVA on diurnal temperature and VPD ranges	20
Table 5:	Wind speed results.....	21
Table 6:	Tests of between-subject effects for MANOVA on relative air temperature and relative VPD.....	24

Abbreviations Used In This Report

ANCOVA	analysis of covariance
DBH	diameter of tree at breast height
MANOVA	multivariate analysis of variance
PAR	photosynthetically active radiation
SE	standard error of the mean
VPD	vapour pressure deficit

Glossary

- Neotropical rainforest** Rainforest located in tropical regions of North and South America.
- Sealed edge** A forest edge with continuous foliage between the forest floor and the forest canopy. This foliage may comprise lateral branches from trees, crowns of saplings and shrubs as well as lianas and seedlings. Sealed edges often develop over time, as foliage growth is stimulated by higher light levels near the forest edge.
- Open edge** Forest edge without continuous foliage between the forest floor and the forest canopy. Forest edges are 'open' when first created and may become sealed over time. Alternatively, forest edges may remain open if other factors prevent them from developing a sealed edge.
- Photosynthetically active radiation** Sunlight between 400 and 700 nm in wavelength, required by plants for photosynthesis.
- Tree successional status** Approximation of the response of a tree species to both disturbance and light availability. Tree species have been classified as either early successional, mid successional or late successional in this report. Early successional species are unable to germinate in shade (i.e. beneath an intact forest canopy), have fast growth rates, require high light levels to survive and grow and are favoured by larger forest disturbances. Late successional species are able to germinate in shade, have seedlings which are able to survive in the shade, have slow growth rates and may persist as suppressed seedlings or saplings for long periods of time beneath the forest canopy. Mid successional species have properties intermediate between early and late successional species. These successional categories also correspond approximately to the positions occupied by species within a sequence of secondary forest succession that occurs when an area of primary forest is cleared.
- Understorey microclimate** Climatic conditions experienced within approximately two metres of the forest floor; includes, but is not limited to, light level, air temperature and humidity, wind speed, soil temperature and soil moisture.
- Vapour pressure deficit** The difference between the moisture holding capacity of saturated air and the actual air moisture. As the air within a leaf's pores is saturated while the air outside the leaf is not, vapour pressure deficit is also a measure of the moisture stress exerted on a plant leaf.

Acknowledgements

Financial support for this project was provided by Powerlink Queensland and the Australian Government's Marine and Tropical Sciences Research Facility. Financial support for pre-cyclone work was provided by Powerlink Queensland, the Rainforest Cooperative Research Centre and the School of Tropical Environmental Studies and Geography, James Cook University.

Mr Gregory Dawe provided assistance with post-cyclone field work. Assistance with pre-cyclone field work was provided by Mrs June Pohlman, Mr Stephen Pohlman, Ms Anne Pohlman, Ms Greta Kading, Mr Nigel Young, Mrs Robin Spencer, Mr Shane Garozzo, Mr Neil Maver, Mr Alex Vaux, Ms Mel Antony, Ms Valery Lopez and Mr Trevor Wilson.

Mr Rigel Jensen identified plant species for the original vegetation survey and Ms Tina Lawson (Research Officer, School of Earth and Environmental Sciences, James Cook University) prepared the maps featured in this report with data from Geosciences Australia.

Field work was conducted under Queensland Environmental Protection Agency scientific research permits WISP00673802 and WITK00672502 and Queensland State Forests permit ATH 02/015.

Introduction

Natural and anthropogenic disturbance near linear canopy openings

Interactions between natural and anthropogenic disturbance may affect the physical structure and species composition of plant communities (Chazdon 2003). Forest fragmentation as a result of human activities has the potential to alter the ecological responses of remaining areas of forest to natural disturbances such as cyclones, fires or droughts (Laurance 1991, 1997; Laurance *et al.* 1997, 2001, 2002; Cochrane and Laurance 2002). In particular, trees at the forest edge may be exposed to greater levels of physical stress and wind damage than trees within continuous forest (McNaughton 1989). Elevated levels of tree damage and mortality have been reported near the edges of tropical forest fragments in central Amazonia, apparently in response to a combination of increased moisture stress and greater wind turbulence (Ferreira and Laurance 1997; Laurance *et al.* 1997, 1998, 2000, 2002; D'Angelo *et al.* 2004). Similarly, elevated levels of wind throw and tree mortality near forest edges have been reported in boreal forests (Essen 1994; Harper *et al.* 2005), temperate forests (Chen *et al.* 1992; Burton 2002; Baldwin and Bradfield 2005; Harper *et al.* 2005) and neotropical rainforests (Williams-Linera 1990; Leigh *et al.* 1993), although other studies have reported no increase in wind throw (Lin *et al.* 2004) or decreased wind throw (Ryall and Smith 2005) near forest or plantation edges. There have been relatively fewer studies of the effects of cyclones and intense storms on vegetation damage in fragmented forests, although elevated levels of cyclone damage to trees have been reported for rainforest edges in northeastern Australia (Laurance 1991, 1997).

Even less is known about the effects of linear infrastructure clearings on the disturbance regimes and ecology of adjacent forest. Tree mortality may be elevated near the edges of such clearings in tropical rainforest in non-cyclonic conditions (Pohlman 2006). However, it is not known whether increased wind exposure at the edges of linear clearings increases rates of vegetation damage and mortality in either normal weather conditions or during intense storms and tropical cyclones. Although there have been anecdotal reports of increased cyclone damage to rainforest near roads and railways in northeastern Australia (Webb 1958), the relationship between proximity to linear clearings and cyclone damage has never been directly examined. If severe wind storms do lead to elevated levels of tree damage and mortality near powerline edges, this may have implications for the dynamics (recruitment, growth and mortality rates [Laurance 1997; Laurance *et al.* 2002]) of rainforest tree communities and subsequently, their species composition and diversity (e.g. Leigh *et al.* 1993; Asquith and Mejia-Chang 2005).

Severe Tropical Cyclone Larry

On 20 March 2006, Severe Tropical Cyclone *Larry* passed directly over the Wooroonooran National Park in the Wet Tropics World Heritage Area of northeastern Queensland. Research sites had previously been established along the Kareeya to Innisfail powerline, along the Palmerston Highway and along Henrietta Creek; all were within the path of the most destructive core of the cyclone (Turton in prep). Initial reconnaissance indicated that these sites were severely damaged (although not completely devastated) and that most of the original site markers had survived the cyclone. This provided a rare opportunity to compare measurements taken before the cyclone with those taken after the cyclone, and to determine whether severe storm damage is greater near the edges of powerline clearings (as well as highway clearings and natural watercourses) than in the rainforest interior.

Research Questions

Two research questions were investigated:

1. Is cyclone damage to trees and saplings greater near the edges of linear canopy openings (powerline clearings, highways and watercourses) than in the rainforest interior?
2. What are the subsequent changes in understorey microclimate (light intensity, air temperature and humidity, soil temperature and moisture and wind speed) near the edges of linear canopy openings?

Methodology

Study Sites

Edge gradients in vegetation damage and microclimate were investigated near three types of linear canopy openings: powerlines, highways and creeks (Figure 1, Figure 2). Field work was conducted in Wooroonooran National Park and South Johnstone State Forest (State Forest 756), in the region around 17° 36' S, 145° 45' E in the Wet Tropics World Heritage Area. Sites were located at elevations between 350 and 720 m above sea level on fertile soils derived from basalt. Prior to Cyclone *Larry*, the forest in this area was complex mesophyll vine forest (Type 1a, *sensu* Tracey 1982). This area falls between the 3000 and 4000 mm annual precipitation isohyets (Siegenthaler and Turton 2000) and the nearest weather station, the South Johnstone Experimental Station (17° 36' 20.16" S, 145° 59' 48.84" E, 18.3m asl) receives an average annual rainfall of 3307 mm, an average daily maximum temperature of 28.1°C and an average daily minimum temperature of 19.1°C (Bureau of Meteorology, www.bom.gov.au). The area experiences a wet season between December and May, with a peak of rainfall between January and March. Forest in this area has been disturbed by selective logging prior to the 1950s and scattered small-scale mining activities prior to the 1990s (Siegenthaler and Turton 2000).

Two sites separated by at least three kilometres were established per edge type (powerline, highway or creek). Two transects, each running for 100 m adjacent to the forest edge were established on opposite sides of the linear canopy opening at each site. Previous studies of rainforests in northeast Queensland have demonstrated that edge gradients in microclimate and vegetation do not extend beyond 25-30 m; thus at 100 m from the edge, conditions are considered to be equivalent to the forest interior (Siegenthaler and Turton 2000; Turton and Freiburger 1997). Due to terrain limitations, it was not possible to place transects exactly opposite each other at each site and in some cases transects were offset by a distance of between 500 m and 3 km. In the case of the second creek site, transects were placed on the same side of the creek but were separated by over 400 m and were established on different orientations (Figure 1). To avoid confounding the effects of edge type with other environmental gradients (e.g. soil type, forest type, annual precipitation), only one linear feature per canopy opening type (powerline versus highway versus creek) was included in the study (the Kareeya to Innisfail powerline, the Palmerston Highway and Henrietta Creek). However, as the sites covered a range of elevations and were physically separated by several kilometres, these data allowed us to assess variation in the nature of the associated edge effects among the three linear canopy opening types within the study area. Full site descriptions are given in Pohlman *et al.* (2007). All field sites were within the area affected by the very destructive core of Tropical Cyclone *Larry* (Turton in prep).




Figure 1: Study area in Wooroonooran National Park and South Johnstone State Forest. Inset P1 shows the layout of powerline site 1 and inset S2 shows the layout of creek site 2 and highway site 2, transect 2. Map produced by Tina Lawson with data from Geosciences Australia (after Pohlman et al. 2007).



Figure 2: Examples of linear canopy openings, prior to Cyclone *Larry*. Clockwise from top left: Palmerston powerline (first three images); Palmerston Highway (following two images); Henrietta Creek; Palmerston Highway.

Vegetation Damage

Vegetation was surveyed at each transect between August 2003 and July 2005 as part of a study on the effects of linear canopy openings on vegetation structure and composition (Pohlman 2006). Trees (stems ≥ 5 cm diameter at breast height [DBH]) were surveyed at 0 m, 4 m, 12 m, 25 m, 50 m and 100 m from the forest edge, within 50 m \times 1 m survey plots running parallel to the forest edge. Saplings (stems 2-4.9 cm DBH) were surveyed within 50 m \times 0.5 m subplots (placed within each tree survey plot). The diameter of all stems was measured and all trees and saplings were identified to species level by a local expert (Mr Rigel Jensen). There were no significant disturbances to the rainforest canopy within the initial survey period.

Six months after Cyclone *Larry* (7-21 September 2006), we revisited these survey plots and were able to determine the fate of 97.67% of stems ≥ 5 cm DBH and 95.38% of stems 2-4.9 cm DBH, as these stems were tagged with brightly-coloured flagging tape which had survived the passage of the cyclone. As leaf resprouting and epicormic growth (*pers. obs.*) had begun by this time, we were unable to assess leaf damage. We recorded damage to branches and stems for each tree and sapling and divided trees and saplings into damage categories of increasing severity (Table 1).

To examine whether vegetation damage was elevated near the edges of linear canopy openings relative to the forest interior, we analysed the data for trees (stems ≥ 5 cm DBH) using an analysis of covariance (ANCOVA) with fixed factors edge type (powerline, highway or creek) and distance from the forest edge (0, 4, 12, 25, 50 and 100 m) and the covariates tree size (diameter at breast height [DBH]), successional status (early-successional ['pioneer' species], mid-successional and late-successional ['climax' or 'primary forest' species]), wood density and growth habit (understorey tree species and canopy tree species). The dependent variable was the damage category recorded for each individual stem. Tree ferns and lianas were not included in this analysis. Data on the growth habit and successional status of each species were obtained from the literature (firstly Cooper and Cooper 1994; Hyland *et al.* 2003; subsequently from Williams 1984, 1987; Osunkoya 1996; Tucker and Murphy 1997; Jackes 2001; Tucker 2001; White *et al.* 2004) and from expert advice (Dr Steve Goosem *pers. comm.*). Data on the wood density of individual tree species were obtained from Osunkoya (1996), Cause *et al.* (1989), Hyland (1989) and from expert advice (Dr Steve Goosem, *pers. comm.*, citing Floyd [1989] and Watson [1951]). Where the wood density of an individual species could not be obtained, a family average (obtained from data in Cause *et al.* 1989 and Smith *et al.* 1991) was used as a surrogate (Appendix 1). In the case of one species (*Phaleria clerodendron*, Thymelaeaceae), no wood density data was available for either the species or the family so the mean wood density of the entire dataset (Appendix 1) was used.

Data on saplings (stems 2-4.9 cm DBH) were analysed using Kruskal-Wallis tests to test for the effects of edge type (powerline, highway or creek) and distance from the edge, as well as growth habit (shrub species, understorey tree species and canopy tree species) and successional status (early-successional, mid-successional and late-successional species) on the damage category recorded for each individual stem. All statistical analyses were performed using SPSS version 11.0.4 for Macintosh.

Table 1: Damage categories for trees and saplings.

Trees (≥ 5 cm DBH)			Saplings (2-4.9 cm DBH)		
Code	Name	Description	Code	Name	Description
0	Intact	No obvious damage.	0	Intact	No obvious damage.
1	Minor branch damage	Few (< 10%) branches lost or only light branch damage sustained.	1	Minor branch damage	Few (< 10%) branches lost or only light branch damage sustained.
2	Bent	Trunk bent < 45° and unbroken.	2	Bent	Trunk bent < 45° and unbroken, tree crown clear of ground and debris.
3	Moderate branch loss	$\leq 50\%$ branches lost or heavily damaged.	3	Pinned	Trunk bent but not broken or uprooted. Substantial part of sapling beneath other debris.
4	Major branch loss	> 50% branches lost or heavily damaged.	4	Moderate branch loss	$\leq 50\%$ branches lost or heavily damaged.
5	Snapped > 10 m	Trunk snapped > 10 m height.	5	Major branch loss	> 50% branches lost or heavily damaged.
6	Snapped 2-10 m	Trunk snapped between 2 and 10 m height.	6	Snapped > 2 m	Trunk snapped > 2 m height.
7	Snapped < 2 m	Trunk snapped < 2 m height [‡] .	7	Snapped < 2 m	Trunk snapped < 2 m height.
8	Uprooted	Visible signs of uprooting [†] .	8	Uprooted	Visible signs of uprooting [†] .

[†] Trees and saplings which were washed out near creek banks were included in category 8 ('uprooted').

[‡] Trees which were flattened beneath tree-fall debris were included in category 7 ('snapped < 2 m').

Understorey Microclimate

We used two complementary methods to investigate the nature of the relationship between understorey microclimate and distance from the forest edge and to explore the effect of vegetation damage from Cyclone *Larry* on these microclimatic edge gradients. Traverse measurements were used to assess the extent and severity of edge gradients in light intensity, soil temperature and soil moisture during daylight hours and data loggers were used to measure diurnal changes in microclimatic edge gradients in air temperature, vapour pressure deficit and wind speed.

Traverse Measurements

The traverse method allows the rapid measurement of spatial variation in forest understorey microclimate (Turton and Freiburger 1997). Measurements began at the outermost point of each transect and continued inwards, towards the forest interior, as swiftly as possible. Measurements were made at 0, 2, 4, 8, 12, 16, 20, 25, 30, 50 and 100 m for all transects (and at -10 m on powerline transects). Previous studies in rainforests in northeast Queensland have demonstrated that edge gradients in microclimate do not extend beyond 25-30 m; thus at 100 m from the edge, microclimatic conditions are considered to be equivalent to forest interior conditions (Turton and Freiburger 1997; Siegenthaler and Turton 2000).

Microclimate measurements were undertaken in clear or relatively clear (cloudy or overcast but not raining) weather, between 10:00 am and 3:00 pm, to avoid the collapse of edge gradients that occurs at low solar angles (Chen *et al.* 1995; Turton and Freiburger 1997).

Pre-cyclone measurements were made in the dry season of 2004 (September 2004; described in Pohlman *et al.* 2007). Pre-cyclone measurements included an expanded range of microclimatic parameters (Pohlman *et al.* 2007) and it took between three and five minutes to take all measurements at any one measuring point and between sixty and ninety minutes to complete a full transect. Post-cyclone measurements were made in the dry season of 2006 (5-6 October 2006), using a reduced set of parameters (the complete set of parameters could not be measured, as damage to the vegetation made it logistically impossible to carry all of the necessary equipment through the forest) and it took approximately 20-25 minutes to complete each transect.

Parameters measured were photosynthetically active radiation (PAR), soil temperature and soil moisture. Photosynthetically active radiation (PAR) (electromagnetic radiation between 400 and 700 nm in wavelength) was measured at a height of 165 cm, using a LiCor quantum sensor (LI-190SA) and meter (LiCor Light Meter, LI-250, Li-COR, Lincoln, Nebraska, USA), which measured the average value of PAR ($\mu\text{mol m}^{-2} \text{sec}^{-1}$) at each point over fifteen seconds. The quantum sensor and display meter were mounted on a 165 cm-tall portable frame of PVC pipe (Figure 3). PAR was measured without simultaneous “open” controls because we were not attempting to measure canopy transmittance; we were interested instead in the mean edge gradients in light level and air speed.

Soil temperature was measured using a 5 cm soil probe (Testo 106-T1, Testo Inc. Flanders, NJ, USA). Pre-cyclone soil moisture (mL/g soil dry weight) was measured from samples taken using a bulb planter to extract soil samples to 10 cm depth. Samples were roughly homogenised and sub-sampled, then weighed prior to and after drying in an oven for a minimum of 48 hours at a temperature of 106°C (until constant weight had been achieved). Post-cyclone soil moisture was measured using a HydroSense soil moisture probe (HydroSense Soil Water Content Measurement System, Campbell Scientific Inc. Logan, UT, USA). Soil moisture measurements were transformed to allow gradients in soil moisture to be compared between years. Proportional soil moisture values were calculated using the following formula:

$$\text{Proportional Value}_{(i)} = \frac{\text{Absolute Value}_{(i)}}{\text{Absolute Value}_{(100\text{m})}} \quad (\text{Eq. 1})$$

for each distance i .

Thus, the value for each of these parameters at the distance of 100m (forest interior) was always 1.

Variations in absolute values of temperature and VPD according to time of day, season and elevation (Chen *et al.* 1995, Turton and Freiburger 1997, Newmark 2001) may obscure edge gradients in these parameters. As the edge gradients are the main focus of interest in this study, this variation was accounted for by transforming the temperature and VPD parameters before analysis. Relative values of soil temperature (this section), air temperature and VPD (next section) were calculated using the following formula:

$$\text{Relative Value}_{(i)} = \text{Absolute Value}_{(i)} - \text{Absolute Value}_{(100\text{m})} \quad (\text{Eq. 2})$$

for each distance i .

Thus, the value for each of these parameters at the distance of 100m (forest interior) was always 0.



Figure 3: The instruments used for the traverse microclimate measurements were supported on a PVC frame. The quantum sensor was attached to the top of the frame (on a levelling mount). The person pictured is Stephen Pohlman. Photo by Catherine Pohlman.

Data for each parameter were analysed using analysis of covariance (ANCOVA) with fixed factors of edge type (powerline, highway and creek) and year (pre-cyclone [2004] and post-cyclone [2006]) and covariate of distance. Distance was included as a covariate, as it is a continuous variable that cannot be broken into discrete treatment levels. Data for PAR were \log_{10} -transformed to approximate normality. Measurements taken at -10m along powerline transects were not included in the analyses, as these values introduced significant outliers into the dataset; values measured at -10 m are shown in the results section for comparison only. Transects were treated as replicates in these analyses ($N = 4$ transects per edge type). Each transect had a different orientation and thus a different interaction with solar angle effects (Turton and Frieberger 1997) and several were also separated by some distance (Figure 1). All statistical analyses were performed using SPSS version 11.0.4 for Macintosh.

Diurnal Variation

Data loggers were used to measure diurnal variation in microclimatic edge gradients. The microclimatic parameters measured were temperature, relative humidity and wind speed. Vapour pressure deficit was calculated from temperature and relative humidity (Jones 1992). Data loggers were placed at distances of 0 m, 4 m, 12 m, 25 m, 50 m and 100 m from the forest edge. For powerline transects, an additional data logger was placed at -10 m from the forest edge. Budget constraints dictated the monitoring of one transect at a time with the data loggers available. Transects were monitored in a semi-random order for a period of at least four consecutive days in each season in which measurements were made.

Air temperature and relative humidity were measured using Tinytag Ultra Temperature and Relative Humidity data loggers (model number TGU-1500) and wind speed was measured using Tinytalk Wind Speed data loggers and cup anemometers (model number TGPR-1201) (Hastings Data Loggers, Port Macquarie, Australia). These instruments were attached to a PVC frame that could be slotted over a wooden stake hammered into the ground (using a spirit level to ensure the stake was placed vertically), in such a way that the TGU-1500 and the cup anemometer of the TGPR-1201 were held at approximately 1.7 m from the ground surface. All external cables had to be protected from the attentions of White-tailed Rats (*Uromys caudimaculatus*) and were housed within thick PVC pipe and multiple layers of electrical tape (Figure 4).

Post-cyclone measurements were made in the dry season of 2006 (between 22 August 2006 and 13 January 2007) and compared with pre-cyclone measurements taken in the dry season of 2004 (between 12 September and 22 December 2004). As there was variation in the length of time data loggers were left at each transect (due to weather and logistical constraints), a subset of four consecutive days of data (the minimum period for which each transect was monitored) was selected from the available data for each transect for each year (the four days selected were either the first four days recorded for each transect or the first four days recorded without large excessive humidity).

To examine the effects of cyclone disturbance and proximity to the forest edge on diurnal variation in air temperature and vapour pressure deficit (VPD), we examined the diurnal range for these parameters for each distance along each transect in each year, by subtracting the overnight minimum value from the following daylight maximum value. This provided a total of three diurnal range values for each distance for each transect, for each year ($n = 3 \text{ days} * 3 \text{ edge types} * 2 \text{ sites} * 2 \text{ transects per site} * 2 \text{ years} = 72$) for both air temperature and VPD. As air temperature and VPD were highly correlated with each other (Pearson correlation $r = 0.931$, $P < 0.001$), we analysed this data using a MANOVA, with the fixed factors of edge type (powerline, highway and creek), distance (0 m, 4 m, 12 m, 25 m, 50 m and 100 m) and year (2004 [pre-cyclone] and 2006 [post-cyclone]). Replication was provided by sites, transects and days. Air temperature and VPD were $\ln(1 + x)$ -transformed prior to analysis to approximate normality. The distance of -10 m measured within the powerline was not included in this analysis, as it would have unbalanced the dataset and introduced significant outliers. The data collected from this point is shown in the results section for comparison only.

As the minimum value for wind speed was 0 km hr^{-1} in all cases, we used the maximum wind speed measured at each distance for each transect over the four days' data collection period for each year as a substitute for "range" ($n = 3 \text{ edge types} * 2 \text{ sites} * 2 \text{ transects per site} * 2 \text{ years} = 24$). Maximum wind speed was also highly correlated with the percent of time for which wind speeds $> 0 \text{ km hr}^{-1}$ were recorded (Spearman rank correlation -0.916 , $P < 0.001$) and so provides a reasonable indication of the overall wind regime at each measuring point. However, this dataset contained a large number of zero values, which skewed the distribution of the data. The data could not be transformed to achieve normality, so non-parametric Kruskal-Wallis tests were used to examine the effects of edge type (powerline, highway and creek), distance (0, 4, 12, 25, 50 and 100 m) and year (pre-cyclone [2004] and post-cyclone [2006]) on maximum wind speed. Measurements at -10 m on powerline transects were not included in this analysis and are shown in the results section for comparison only.

To complement the instantaneous measurements of PAR, soil temperature and soil moisture taken using the traverse method (see previous section), gradients in air temperature and VPD were calculated for three periods within the day: morning (10:00-10:50), midday (12:00-12:50) and afternoon (14:00-14:50). Relative air temperature and relative VPD were calculated for each ten minute reading within these time periods (Eq. 2; see previous section)

and average relative values were calculated for each time period for three days per transect ($N = 12$ transects \times 3 days per transect \times 3 time periods per day = 108). These average hourly values were used as replicates in a MANOVA, with fixed factors of edge type (powerline, highway and creek), distance (0, 4, 12, 25 and 50 [values for 100 m were always 0 and were excluded from the analysis to allow the dataset to approximate normality]) and time of day (morning [10:00-10:50], midday [12:00-12:50] and afternoon [14:00-14:50]). Data for 2004 (pre-cyclone) and 2006 (post-cyclone) were analysed separately, as the combined dataset could not be transformed to approximate normality. All statistical analyses were performed using SPSS version 11.0.4 for Macintosh.



Figure 4: The instruments used to measure diurnal variation in microclimate were supported on a PVC frame resting on a wooden post. Instruments were attached at a height of 170 cm above the ground surface. The person pictured is Stephen Pohlman, who is standing behind and slightly upslope of the apparatus. Photo by Catherine Pohlman.

Results

Vegetation Damage

Tree Damage

Tree damage was influenced by size (diameter at breast height [DBH]), successional status and wood density (Table 2). Early-successional tree species (i.e. pioneer tree species, which tend to have fast growth rates, low wood density and to be favoured by high light levels and/or larger disturbances (Turner 2001) suffered greater damage than mid- or late-successional tree species (late-successional tree species [also called ‘climax’ species or ‘primary forest species’], are species able to germinate in shade [i.e. beneath the forest canopy] and which tend to have slower growth rates, long persistence times in the seedling and sapling pools of the forest understorey and higher wood densities [Turner 2001]) (Figure 5a). There were also non-significant trends towards lesser damage with decreasing diameter (Figure 5b) and greater damage with lower wood density (Figure 5c, Table 2). In addition, there was a significant edge type \times distance interaction (Table 2); tree damage was slightly lower near powerline clearing edges (Figure 6a) and slightly higher near creek edges than in the forest interior (Figure 6c) but was not influenced by distance from highway edges (Figure 6b). However, post-hoc tests for the effects of distance within each edge type were not significant ($P > 0.05$).

Table 2: Cyclone damage to trees (stems ≥ 5 cm DBH). Tree damage data were analysed with ANCOVA (covariates DBH, successional status, wood density and growth habit; fixed factors edge type and distance). Significant effects are highlighted in bold and marginally significant effects are highlighted with italics.

Effect	<i>F</i>	d.f.	<i>P</i>
Covariates			
Tree Size (DBH)	2.812	1	<i>0.094</i>
Successional Status	9.586	1	0.002
Wood Density	3.305	1	<i>0.069</i>
Growth Habit	1.004	1	0.316
Fixed Factors			
Edge Type	6.269	2	0.002
Distance	1.863	5	<i>0.098</i>
Edge Type \times Distance	1.830	10	0.052





Figure 5: Covariates of cyclone damage to trees (stems ≥ 5 cm DBH). Damage severity was influenced by (a) successional status ($p = 0.002$), (b) tree size (DBH) ($p = 0.094$) and (c) wood density ($p = 0.069$). Left-hand y-axes indicate the percentage of stems falling within different damage categories and right-hand y-axes indicate mean damage scores.



Sapling Damage

Sapling damage was influenced by a number of factors (Table 3). Of the two covariates tested, sapling damage was related to growth habit but not successional status (Table 3). Sapling damage tended to be greater in species with taller adult stature: sapling damage was greatest in tree species that reach the canopy when fully grown, intermediate in tree species that only occur in the understorey as adults and least in shrub species (Figure 7). Of the main factors examined (edge type and distance from the forest edge), the edge type × distance interaction was significant (Table 3). Sapling damage was lower near the very edge (0 m) of powerline clearings (Figure 8a) and higher near creek edges (Figure 8c) than in the forest interior and elevated between 12 and 25 m from highway edges (Figure 8b).

Table 3: Cyclone damage to saplings (stems 2-5 cm DBH). Sapling damage data were analysed with Kruskal-Wallis tests. Significant effects are highlighted in bold and marginally significant effects are highlighted with italics.

Effect	χ^2	d.f.	<i>P</i>
Covariates			
Growth Habit	13.606	2	0.001
Successional Status	4.253	2	0.119
Fixed Factors			
Edge Type	1.591	2	0.451
Distance	8.059	5	0.153
Edge Type × Distance	37.571	17	0.003





Figure 7: Covariate of cyclone damage to saplings (stems 2-4.9 cm DBH). Cyclone damage varied between species with differing growth habits ($P = 0.001$) Left-hand y-axes indicate the percentage of stems falling within different damage categories and right-hand y-axes indicate mean damage scores.



Damage Category

- Intact
- Minor Branch Loss
- Bent
- Pinned
- Moderate Branch Loss
- Major Branch Loss
- Snapped > 2 m
- Snapped < 2 m
- Uprooted

Edge Type

- Powerline
- Highway
- ▲ Creek

Figure 8: Cyclone damage to saplings (stems 2 – 4.9 cm DBH). Variation in damage severity with distance is shown for powerline (a), highway (b) and creek (c) transects. Left-hand y-axes indicate the percentage of stems falling within different damage categories and right-hand y-axes indicate mean damage scores. Symbols which share letters were not significantly different ($P > 0.05$, *post-hoc* Mann-Whitney *U* tests).

Understorey Microclimate

Traverse Measurements

Vegetation damage caused by Severe Tropical Cyclone *Larry* affected the direction and severity of microclimatic edge gradients measured near the edges of powerlines, highways and streams. Photosynthetically active radiation (PAR) was bimodally distributed; values of PAR measured six months after Cyclone *Larry* were much greater than those measured before the cyclone ($t = 31.468$, $P < 0.001$). Data for PAR for 2004 (pre-cyclone) and 2006 (post-cyclone) were thus analysed separately. Prior to the cyclone, there was a strong increase in light intensity near the edges of all three types of linear canopy openings (covariate distance, $F = 11.581$, d.f. = 1, $P = 0.001$; Figure 9 a, b, c), although creek edges had greater light availability than powerline or highway edges (main effect edge type, $F = 7.889$, d.f. = 2, $P = 0.001$). After the cyclone, the edge gradients in light intensity had disappeared (covariate distance, $F = 0.814$, d.f. = 1, $P = 0.369$; Figure 9), although light availability was still marginally higher along creek transects than along powerline or highway transects (main effect edge type, $F = 2.783$, d.f. = 2, $P = 0.066$).




Figure 9: Photosynthetically active radiation (PAR) for (a) powerline, (b) highway and (c) creek transects, measured in the dry seasons eighteen months before (open symbols) and six months after (closed symbols) Cyclone *Larry*. Note the log-scale on the y-axis. Dashed vertical lines indicate the position of the forest edge. Values represent means ± 1 SE.

Gradients in soil temperature differed among edge types and between years (edge type \times year interaction, $F = 12.743$, d.f. = 2, $P < 0.001$; Figure 10). Although the effect of distance on relative soil temperature was not significant (covariate distance, $F = 2.871$, d.f. = 1, $P = 0.091$), some trends were apparent. Near powerline edges, soil temperatures tended to be slightly lower near the very edge of the forest (within ~ 4 m of the edge) than in the forest interior and this pattern was only slightly altered after the cyclone (Figure 10a), possibly due to a slightly higher level of vegetation damage within the forest interior (Figure 6a) exposing the forest to greater light availability and harsher microclimate. Near highway edges, soil

temperatures were slightly elevated at the very edge (within ~2 m of the edge) of the forest prior to the cyclone; six months after the cyclone, soil temperatures were lower near the forest edge and elevated at intermediate distances (~ 12-30 m) relative to the forest interior (Figure 10b). Again this may relate to the larger amounts of damage to saplings at these intermediate distances allowing greater light penetration and thus soil temperature to rise. In contrast, near creek edges, soil temperatures were slightly elevated at the very edge of the forest (within ~4 m of the edge) prior to the cyclone, whereas, six months after the cyclone, soil temperatures were greater in the interior of the forest than within 50 m of the edge (Figure 10c).




Figure 10: Relative soil temperature for (a) powerline, (b) highway and (c) creek transects, measured in the dry seasons eighteen months before (open symbols) and six months before (closed symbols) Cyclone *Larry*. Dashed vertical lines indicate the position of the forest edge and dashed horizontal lines indicate the relative value of measurements taken in the forest interior (100 m). Values represent means ± 1 SE.

Soil moisture tended to be lower near the forest edge than in the forest interior (covariate distance, $F = 3.785$, d.f. = 1, $P = 0.053$; Figure 11), although gradients in soil moisture differed among edge types and between years (edge type \times year interaction, $F = 9.496$, d.f. = 2, $P < 0.001$). Prior to the cyclone, soil moisture tended to decrease slightly from powerline edges to the forest interior but six months after the cyclone, this gradient had reversed and soil moisture was slightly lower near powerline edges than in the forest interior (Figure 11a). There was no clear gradient in soil moisture near highway edges prior to the cyclone but soil moisture was slightly lower near the very edge of the forest (~ 0-4 m) six months after the cyclone (Figure 11b). Soil moisture was lower near creek edges than in the forest interior both before and after the cyclone, although soil moisture obtained interior values closer to the edge after the cyclone than before the cyclone (Figure 11c).




Figure 11: Proportional soil moisture for (a) powerline, (b) highway and (c) creek transects, measured in the dry seasons eighteen months before (open symbols) and six months after (closed symbols) Cyclone *Larry*. Dashed vertical lines indicate the position of the forest edge and dashed horizontal lines indicate the relative value of measurements taken in the forest interior (100 m). Values represent means ± 1 SE.

Diurnal Variation

Severe Tropical Cyclone *Larry* altered both the absolute values of and the edge gradients in the diurnal ranges of air temperature and vapour pressure deficit (VPD). The direction of edge gradients differed between years (distance \times year interaction; Pillai's trace = 0.090, $F = 3.521$, d.f. = 10, $P < 0.001$); in addition, diurnal ranges of air temperature and VPD differed among edge types and between years (edge type \times year interaction; Pillai's trace = 0.051, $F = 4.820$, d.f. = 4, $P = 0.001$), although the three-way interaction was not significant (edge type \times distance \times year interaction; Pillai's trace = 0.019, $F = 0.359$, d.f. = 0.359, d.f. = 20, $P = 0.996$). Prior to the cyclone, diurnal ranges of air temperature and VPD were slightly greater near the very edges of powerlines and highways than in the forest interior (Figure 12 a, b, d, e), but did not vary with distance from creek edges (Figure 12 c, f). After the cyclone, these gradients had reversed, and diurnal ranges of air temperature and VPD were lower near the edges of powerlines, highways and streams than in the forest interior. The edge type \times year interaction detected in the MANOVA was not supported by the *post hoc* tests of between-subject effects for the individual parameters (Table 4); neither air temperature nor VPD displayed significant edge type \times year interactions in these tests.

Maximum wind speed generally decreased from the forest edge to the forest interior, for all three edge types (Figure 13 a, b, c; Table 5). Wind speeds were greater along creek transects than along powerline or highway transects, both prior to the cyclone and six months after it. Additionally, although wind speed was greater after the cyclone than before the cyclone, this increase was greatest along creek transects and least along powerline transects (Figure 13, a, b, c; Table 5) and the increase in wind speeds along highway transects was somewhat greater at intermediate distances than near the edge or in the forest interior (Figure 8b). In general, wind speed was greater for distances between 4 and 100 m

from the forest edge after the cyclone than before the cyclone (Figure 13, a, b, c; Table 5), indicating a greater penetration of wind into the forest interior after the cyclone.

Edge gradients in relative air temperature and relative VPD differed between years and among edge types (Table 6). Prior to the cyclone, air temperature and VPD were higher near the forest edge than in the forest interior, but this increase was greater near powerline and highway edges than near stream edges (edge type \times distance interaction; Pillai's trace = 0.132, $F = 4.223$, d.f. = 16, $P < 0.001$; Table 6; Figure 14 a, b, c, g, h, i). In addition, the magnitude of edge gradients increased between morning and afternoon (time of day main effect; Pillai's trace = 0.038, $F = 4.653$, d.f. = 4, $P = 0.001$; Table 6; Figure 14 a, b, c, g, h, i). Six months after the cyclone, edge gradients in air temperature and VPD varied with time of day and among edge types (edge type \times distance \times time of day; Pillai's trace = 0.111, $F = 1.626$, d.f. = 32, $P = 0.016$; Table 6; Figure 14 d, e, f, j, k, l). Along powerline transects, air temperature and VPD were lower near the edge than in the forest interior and this edge gradient was strongest in the morning and, although still present at midday and in the afternoon, intermediate distances displayed higher temperature and VPD compared to the forest interior (Figure 14 d, j). Along highway transects, air temperature and VPD were lower near the forest edge than in the forest interior and this gradient was strongest in the morning and lesser in the afternoon (Figure 14 e, k). Along creek transects, air temperature and VPD were slightly higher near the forest edge (between 4 and 50 m from the edge) than in the forest interior in the morning but were lower near the edge than in the forest interior at midday and in the afternoon, with the exception of the 50 m point at 2:00 pm, which displayed higher values of air temperature and VPD than either the edge or the forest interior (Figure 14 f, l).

Table 4: Tests of between-subject effects (distance, edge type and year) for the MANOVA on diurnal temperature ($^{\circ}\text{C}$) and VPD (kPa) ranges. Significant effects have been highlighted in bold.

Factor	Parameter	<i>F</i>	d.f.	<i>P</i>
Edge Type	<i>ln</i> (1 + diurnal temperature range)	0.617	2	0.540
	<i>ln</i> (1 + diurnal VPD range)	1.607	2	0.202
Distance	<i>ln</i> (1 + diurnal temperature range)	1.767	5	0.119
	<i>ln</i> (1 + diurnal VPD range)	1.769	5	0.118
Year	<i>ln</i> (1 + diurnal temp)	292.747	1	< 0.001
	<i>ln</i> (1 + diurnal VPD range)	264.864	1	< 0.001
Edge Type \times Distance	<i>ln</i> (1 + diurnal temperature range)	0.896	10	0.537
	<i>ln</i> (1 + diurnal VPD range)	0.417	10	0.939
Edge Type \times Year	<i>ln</i> (1 + diurnal temperature range)	2.632	2	0.073
	<i>ln</i> (1 + diurnal VPD range)	1.557	2	0.212
Distance \times Year	<i>ln</i> (1 + diurnal temperature range)	5.877	5	< 0.001
	<i>ln</i> (1 + diurnal VPD range)	5.864	5	< 0.001
Edge Type \times Distance \times Year	<i>ln</i> (1 + diurnal temperature range)	0.544	10	0.859
	<i>ln</i> (1 + diurnal VPD range)	0.547	10	0.856

Table 5: Wind speed (km hr^{-1}) results table. Each effect analysed with Kruskal-Wallis tests (or Mann-Whitney U tests for ‘Year’). Significant effects are highlighted in bold.

Effect	χ^2	d.f.	<i>P</i>
Edge Type	18.602	2	< 0.001
Distance	35.534	5	< 0.001
Year	1684.000 [†]	1	0.003
Edge Type × Distance	59.902	17	< 0.001
Edge Type × Year	30.004	5	< 0.001
Distance × Year	52.728	11	< 0.001
Edge Type × Distance × Year	81.712	35	< 0.001

[†] Mann-Whitney U test.




Figure 12: Diurnal temperature range (a, b, c) and diurnal vapour pressure deficit (VPD) range (d, e, f) for powerline (a, d), highway (b, e) and creek (c, f) transects in the dry seasons eighteen months before (open symbols) and six months after (closed symbols) Cyclone *Larry*. Dashed vertical lines indicate the position of the forest edge. Values represent means ± 1 SE.




Figure 13: Maximum wind speed for powerline (a), highway (b) and creek (c) transects in the dry seasons eighteen months before (open symbols) and six months after (closed symbols) Severe Tropical Cyclone *Larry*. Dashed vertical lines indicate the position of the forest edge. Values represent means ± 1 SE.




Figure 14: Edge gradients in air temperature and VPD in the morning (10:00-10:50), at midday (12:00-12:50) and in the afternoon (14:00-14:50). Dashed vertical lines indicate the position of the forest edge and dashed horizontal lines indicate the relative value of measurements made in the forest interior (100 m). Values represent means \pm 1 SE.

Table 6: Tests of between-subject effects (distance, edge type and time of day) for the MANOVA on relative air temperature (°C) and relative VPD (kPa). Data for 2004 (pre-cyclone) and 2006 (post-cyclone) were analysed separately. Significant effects ($\alpha = 0.05$) are highlighted in bold.

Effect	Parameter	Pre-cyclone (2004)			Post-cyclone (2006)		
		<i>F</i>	d.f.	<i>P</i>	<i>F</i>	d.f.	<i>P</i>
Edge Type	Relative Temperature (°C)	4.136	2	0.017	0.298	2	0.742
	Relative Vapour Pressure Deficit (kPa)	1.441	2	0.238	0.180	2	0.835
Distance	Relative Temperature (°C)	47.035	4	< 0.001	7.131	4	< 0.001
	Relative Vapour Pressure Deficit (kPa)	39.206	4	< 0.001	8.983	4	< 0.001
Time of Day	Relative Temperature (°C)	3.845	2	0.022	2.272	2	0.104
	Relative Vapour Pressure Deficit (kPa)	8.882	2	< 0.001	2.768	2	0.064
Edge Type × Distance	Relative Temperature (°C)	6.874	8	< 0.001	0.820	8	0.585
	Relative Vapour Pressure Deficit (kPa)	5.006	8	< 0.001	1.209	8	0.292
Edge Type × Time of Day	Relative Temperature (°C)	1.122	4	0.345	6.614	4	< 0.001
	Relative Vapour Pressure Deficit (kPa)	1.201	4	0.310	5.111	4	< 0.001
Distance × Time of Day	Relative Temperature (°C)	0.743	8	0.653	1.773	8	0.080
	Relative Vapour Pressure Deficit (kPa)	0.707	8	0.685	1.651	8	0.108
Edge Type × Distance × Time of Day	Relative Temperature (°C)	0.696	16	0.799	2.565	16	0.001
	Relative Vapour Pressure Deficit (kPa)	0.559	16	0.914	2.602	16	0.001

Discussion

Vegetation Damage

Effect of proximity to linear clearings

Damage to trees was slightly reduced near powerline edges and slightly elevated near creek edges, relative to the forest interior (Figure 6). Similarly, sapling damage was reduced near powerline edges and elevated near creek edges, relative to the forest interior but was also elevated at intermediate distances from highway edges (Figure 8). These results contradicted our initial hypothesis that vegetation damage from Cyclone *Larry* would be generally greater near the edges of linear canopy openings than in the forest interior; vegetation damage was elevated near creek edges, but not near anthropogenic edges (powerlines and highways). In light of the literature documenting greater wind damage near forest edges in tropical, temperate and boreal forests (Williams-Linera 1990; Laurance 1991, 1997; Chen *et al.* 1992; Leigh *et al.* 1993; Essen 1994; Ferreira and Laurance 1997; Laurance *et al.* 1997, 1998, 2000, 2002; Burton 2002; D'Angelo *et al.* 2004; Baldwin and Bradfield 2005; Harper *et al.* 2005), this result is surprising, although such patterns are not entirely ubiquitous (e.g. Lin *et al.* 2004; Ryall and Smith 2005). Few of these studies, however, have specifically examined the damage caused by severe wind storms.

The severe winds associated with Cyclone *Larry* (gusts to 294 km hr⁻¹ at Bellenden Ker [Bureau of Meteorology 2007]) appear to have caused vegetation destruction so severe that it overrode any smaller-scale landscape effects (Everham and Brokaw 1996), although effects caused by larger-scale topographic features were evident (Turton in prep). Although elevated cyclone damage has previously been detected near forest edges bordering agricultural land on the Atherton tablelands (approximately 20 km west of our study sites) (Laurance 1991, 1997), this damage was caused by a category 3 cyclone, with maximum wind gusts of 176 km hr⁻¹ (Unwin *et al.* 1988), which caused less severe vegetation damage in the study area than Cyclone *Larry* (Unwin *et al.* 1988; Turton in prep). If forest edges do influence the vulnerability of vegetation to cyclone damage, this effect may be confined to areas of less severe overall cyclone damage. A study of broader landscape patterns associated with Cyclone *Larry* is currently underway. This project, also funded by Powerlink, is examining cyclone damage over a wider area of the Palmerston region using remote sensing techniques applied to satellite and aerial photographic imagery to examine tree fall severity and direction and vegetation indices (Goosem *et al.* in prep.).

Effect of tree size and successional status

Tree damage was more severe for early successional tree species. This is consistent with the trend towards increasing damage with decreasing wood density, as early successional species tend to have lower wood densities (Osunkoya 1996; Turner 2001; Falster and Westoby 2005; van Gelder *et al.* 2006; Osunkoya *et al.* 2007). Tree damage also tended to be more severe in larger trees than in smaller trees. Damage to saplings was greater for individuals from taller-growing species (canopy trees) than for shorter-growing species (understorey trees and shrubs), although damage did not differ among species from different successional stages. We could not assess the effects of wood density on sapling damage, but wood density tends to be greater in species with shorter stature, at least among shade-tolerant species (Falster and Westoby 2005; van Gelder *et al.* 2006). As wood density is negatively correlated with damage rates (Falster 2006), these data suggest that, unsurprisingly, species with higher wood density (and thus greater resistance to damage under non-cyclonic conditions), suffered lower rates of cyclone damage than species with lower wood density. Similar patterns of greater cyclone damage to species with lower wood

density and/or greater damage to pioneer species have been detected in many previous studies around the world (e.g. Zimmerman *et al.* 1994; Everham and Brokaw 1996; Ostertag *et al.* 2005; Franklin *et al.* 2004), although Grove *et al.* (2000) observed greater cyclone damage in old-growth stands than in second-growth stands in the Daintree area of northeastern Queensland after Tropical Cyclone *Rona*.

Cyclone damage tended to be greater for trees with larger diameters (Figure 5b). This relationship appeared to take the form of a threshold response, rather than a simple linear correlation, with trees less than approximately 20 cm in diameter experiencing lower cyclone damage than trees with larger diameters. However, this relationship was not significant at the $\alpha = 0.05$ level and there was a higher proportion of snapped and uprooted trees in the smaller size classes than in the larger size classes. In contrast, the proportion of trees with significant branch damage was greater amongst larger trees than amongst smaller trees. This pattern of damage suggests that larger trees, which tend to have larger crowns and thus greater crown exposure in the canopy (Poorter *et al.* 2006), suffered greater crown damage but less major stem damage (snapping and uprooting) than smaller trees (with correspondingly smaller crowns and thus lower crown exposure). It is possible that much of the damage suffered by smaller trees was due to falling branches and other debris from the forest canopy, while much of the damage to larger trees was due to the direct effects of strong winds from the cyclone (e.g. Everham and Brokaw 1996).

Effects of 'sealed' edges

The different edge patterns that were observed in the current study (i.e. slightly reduced tree and sapling damage near powerlines, slightly elevated tree and sapling damage near creeks and slightly elevated sapling damage at intermediate distances from highway edges) may also have resulted from differences in pre-cyclone edge conditions among edge types. Prior to Cyclone *Larry*, there were higher proportions of smaller trees and saplings near the forest edge than in the forest interior, for all three edge types, as well as increased abundances of lianas and weeds (Pohlman 2006). Forest near creek edges, however, had a more open canopy structure (which allowed a greater penetration of light and wind into the forest understorey), a greater proportion of pioneer species and a greater abundance of lianas than forest near powerline or highway edges (Pohlman 2006). These differences were ascribed to differences in the degree of edge 'sealing' that occurred in the time since the current forest edges were created.

Edge sealing occurs when the increased light availability at a newly-formed forest edge encourages the growth of foliage on remaining trees, as well as the growth of saplings, seedlings and lianas, until a 'wall of vegetation' covers the forest edge from the ground to the forest canopy (Laurance *et al.* 2002; Harper *et al.* 2005). Sealed edges have been found to decrease the severity of some, though not all, microclimatic and biotic edge effects (e.g. Camargo and Kapos 1995; Kapos *et al.* 1997; Cadenasso and Pickett 2000, 2001; Harper *et al.* 2005). Prior to Cyclone *Larry*, forest edges near powerlines and highways were more sealed than forest edges near creeks. Edges near anthropogenic linear clearings had a higher foliage density (measured indirectly through a decrease in the *red : far red* ratio), reduced recruitment of pioneer species and less dramatic increases in understorey light availability and the abundance of lianas than forest near creek edges (Pohlman 2006). These differences in edge sealing may have been due to differences in the frequency of physical disturbance to the forest edge. Creek edges may be subject to periodic flood disturbance, whilst powerline and highway edges have not been significantly disturbed in approximately fifteen years (although highway edges may receive some disturbance from periodic highway maintenance activities) (Pohlman *et al.* 2007).

The greater proportion of pioneer species and the open edge structure near creeks, as well as the sandier soil found near creek edges (Pohlman *et al.* 2007) may have increased the

vulnerability of trees and saplings to damage from Cyclone *Larry*. Although Cyclone *Larry* was a 'dry' cyclone, in that it was not associated with heavy rainfall (Turton in prep), sections of the creek bank appeared to have been washed away or undermined during post-cyclone floods (*personal observation*), which may have accounted for a large proportion of the greater number of uprooted trees and saplings near creek edges (Figures 6c, 8c). As pioneer species also suffered greater cyclone damage, the relatively elevated abundance of these species near creek edges is also likely to have contributed to the high cyclone damage observed near creek edges. Conversely, the relatively lower abundance of pioneer species and more sealed edge structure near powerline and highway edges may have offered some protection from cyclone damage. A thin strip of edge vegetation with foliage slightly more intact than that in the forest interior was observed near highway edges (*personal observation*; see also Figure 9b, where lowered PAR is evident near highway edges) although not near powerline edges. It is possible that the lower proportion of large trees (which were more vulnerable to cyclone damage than smaller trees) left vegetation at the edge less exposed to damage from falling branches and canopy debris and, in addition, that the relatively low proportion of pioneer species near powerline and highway edges (as compared with creek edges) also decreased the vulnerability of these forest edges.

Understorey Microclimate

Prior to Cyclone *Larry*, edge gradients were present in understorey microclimate. For example, light availability increased towards the forest edge, although this increase was greatest near creek edges (Figure 9). Prior to the cyclone there were only very slight changes in soil temperature at the very edge of the forest (Figure 10) and soil moisture was lower near creek edges (Figure 11c), possibly due to higher sand content (and thus lower moisture holding capacity) of soil near the creek edge (Pohlman *et al.* 2007). However, edge gradients in air temperature and vapour pressure deficit were obvious, being elevated at the forest edge near powerlines and highways but not creeks, while the strength of these gradients increased between morning and afternoon (Figure 14). Additionally, the diurnal ranges of temperature and vapour pressure deficit were greater at the edge of the forest, near powerlines and highways but not creeks (Figure 12). Finally, wind speed increased near the forest edge, although this increase was greatest near creek edges (Figure 13). Following Cyclone *Larry*, the severe damage to the forest canopy led to an increase in light availability and wind speed in the forest understorey (Figures 9, 13) and the loss or alteration of numerous microclimatic edge gradients.

Following the cyclone, edge gradients in light availability were lost (Figure 10), with the exception of a decrease in light availability near highway edges (Figure 10b) that was associated with a thin strip of edge vegetation that was less damaged than the surrounding forest (*personal observation*). Edge gradients in the diurnal ranges of air temperature and vapour pressure deficit were reversed after the cyclone (Figure 12); temperature and VPD were lower near the forest edge than in the forest interior, for all three edge types. Maximum wind speeds were elevated near the forest edge both before and after the cyclone, but both the absolute values of wind speed and the magnitude of the edge gradients in wind speed increased near creek edges and highway edges following the cyclone. Edge gradients in air temperature and vapour pressure deficit observed during daylight hours (Figure 14) became more complex after the cyclone.

Increases in light availability and wind speed in the forest understorey reflect the severe cyclone damage to the rainforest canopy (Turton 1992; Turton and Siegenthaler 2004). The high foliage density of intact rainforest canopies moderates the microclimate experienced in the rainforest understorey, decreasing light intensity and wind speed and increasing humidity (e.g. Ray *et al.* 2005). Prior to the cyclone, light levels measured in the forest interior were approximately < 0.2-7 % of those measured in the open. Six months following the cyclone,

interior light measurements had risen to approximately 3-100 % of open values. This dramatic increase in light availability in the forest understorey was complemented by increases in wind speed and diurnal ranges of air temperature and vapour pressure deficit. In contrast to light availability, increases in wind speed were not accompanied by the loss of edge gradients (Figure 9). The persistence of edge gradients in wind speed indicates that, even after the cyclone, certain edge processes were still operating in the forest understorey. The increase in the absolute values of wind speed reflects the importance of canopy structure for the wind regime experienced in the forest understorey (e.g. Ray *et al.* 2005). Similarly, the increase in wind speeds at intermediate distances from highway edges is consistent with the combined effects of increased tree and sapling damage found at those distances.

Changes in edge gradients of air temperature and vapour pressure deficit (Figures 12 and 14) may reflect the degree of damage to the forest canopy. Although the diurnal edge gradients became more complex after the cyclone (Figure 14), there is a general trend towards a reversal of the edge gradients present before the cyclone. Prior to the cyclone, both the diurnal ranges of temperature and VPD (Figure 12) and the diurnal edge patterns (Figure 14) tended to show positive edge gradients (i.e. values were greater near the edge than in the forest interior) near powerline and highway edges. Following the cyclone, edge gradients became negative (i.e. values were lower near the forest edge than in the forest interior) near powerline and highway edges. It is possible that the slightly lower vegetation damage near powerline edges (Figures 6a and 8a) and the strip of vegetation with relatively intact foliage near highway edges (*personal observation*) provided a slightly more closed canopy (and thus a greater degree of insulation from the prevailing temperature, sunlight and wind conditions) near the forest edge than in the more extensively damaged forest interior. Changes in edge gradients near creeks appeared slightly less pronounced, but were roughly consistent with those measured near powerlines and highways. Given the greater vegetation damage observed near creek edges (Figures 6c and 8c), this pattern is unlikely to be a direct result of differences in canopy damage between the creek edge and the forest interior. However, prior to the cyclone, the lack of edge gradients in air temperature and VPD near creeks was ascribed to a buffering effect of increased evaporative cooling from the flowing water within the creek itself (Pohlman *et al.* 2007) and it is possible that this mechanism is still operating to reduce the extremes of temperature and VPD experienced near the severely damaged creek edge.

Cyclone Larry and Linear Canopy Openings

Contrary to our initial expectations, cyclone damage was not greater near the edges of powerlines and highways than in the forest interior, although cyclone damage was elevated near creek edges. The greater degree of vegetation 'sealing' of forest edges near powerlines and highways, as well as the relatively low abundance of tree species with traits associated with increased vulnerability to wind damage (e.g. low wood density and early successional status) and the greater proportion of trees with smaller diameters near the forest edge, may have offered some degree of protection from the more severe cyclone damage observed near creek edges. Additionally, flood damage near the creek edges probably amplified the wind effects.

The severe canopy damage caused by Cyclone *Larry* led to dramatic changes in the microclimate of the rainforest understorey. Light availability increased dramatically after the cyclone, as did wind speed and the diurnal ranges of air temperature and vapour pressure deficit. Slightly less severe vegetation damage near the edges of powerlines and highways led to slightly reduced temperature and VPD extremes, relative to the forest interior. The increase in light availability since the cyclone has been sufficient to allow the germination and growth of numerous pioneer and weed species (*personal observation*) and it is possible that

the increased wind speeds (and continued presence of edge gradients in wind speed) may facilitate the invasion of weed species into the rainforest (e.g. Bellingham *et al.* 2005), as many weed species of more open habitats are wind-dispersed. Species common in the Palmerston powerline clearing and along the road and highway edges such as Guinea grass (*Urochloa maxima*) and molasses grass (*Melinis minutiflora*) and a variety of herbaceous weeds (Goosem 2006; Pohlman 2006) are wind-dispersed and have been observed colonising the cyclone-disturbed areas (*personal observations*). Recovery of the forest canopy may lead to a decrease in the abundance of pioneer and weed species (e.g. Burslem *et al.* 2000), however, should weed species capable of interfering with canopy recovery become established, this process may be considerably slowed, if not stalled altogether (Gascon *et al.* 2000). In the linear infrastructure clearings of this area, weed species such as the grasses mentioned above, as well as the scramblers *Lantana camara* and *Rubus alceifolius* have a tendency to suppress the regeneration of rainforest tree seedlings (Reynolds 1994; Werren 2001; Goosem and Turton 2006). The post-cyclone situation therefore requires close monitoring.

Recommendations

We make the following recommendations:

1. Where it is not possible to minimise the footprint of linear infrastructure corridors (e.g. to clear only the area around the base of each transmission line tower), a 'sealed' edge structure should be maintained. The sealed edge structure near powerlines and highways appears to have offered some degree of protection from the more severe wind damage observed near creek edges. A sealed edge structure may be encouraged and maintained through measures which minimise the disturbance to the forest edge during infrastructure and easement maintenance (e.g. avoid mechanical disturbance to the forest edge; avoid the use of fire near rainforest edges [e.g. Cochrane and Laurance 2002]).
2. Microclimatic changes following Cyclone *Larry* may favour the germination and growth of numerous pioneer and weed species and initial observations suggest that the abundance of exotic plant species within the rainforest understorey has increased substantially since the cyclone. We recommend that this situation be monitored, in order to identify potential problems (e.g. incursions of scrambling weeds [e.g. *Lantana camara*, *Rubus alceifolius*] or grasses into the forest).

References

- Asquith, N. M. and Mejia-Chang, M. (2005) Mammals, edge effects, and the loss of tropical forest diversity. *Ecology* 86, 379-390.
- Baldwin, L. K. and Bradfield, G. E. (2005) Bryophyte community differences between edge and interior environments in temperate rain-forest fragments of coastal British Columbia. *Canadian Journal of Forest Research* 35, 580-592.
- Bellingham, P. J., Tanner, E. V. J. and Healey, J. R. (2005) Hurricane disturbance accelerates invasion by the alien tree *Pittosporum undulatum* in Jamaican montane rain forests. *Journal of Vegetation Science* 16, 675-684.
- Bureau of Meteorology (2007) *Severe Tropical Cyclone Larry*. Australian Government, Bureau of Meteorology. Available online at www.bom.gov.au/weather/qld/cyclone/tc_larry/
- Burslem, D. F. R. P., Whitmore, T. C. and Brown, G. C. (2000) Short-term effects of cyclone impact and long-term recovery of tropical rain forest on Kolombangara, Solomon Islands. *Journal of Ecology* 88, 1063-1078.
- Burton, P. J. (2002) Effects of clearcut edges on trees in the sub-boreal spruce zone of Northwest-Central British Columbia. *Silva Fennica* 36, 329-352.
- Cadenasso, M. K. and Pickett, S. T. A. (2000) Linking forest edge structure to edge function: mediation of herbivore damage. *Journal of Ecology* 88, 31-44.
- Cadenasso, M. L. and Pickett, S. T. A. (2001) Effect of Edge Structure on the Flux of Species into Forest Interiors. *Conservation Biology* 15, 91-97.
- Camargo, J. L. C. and Kapos, V. (1995) Complex edge effects on soil moisture and microclimate in central Amazonian forest. *Journal of Tropical Ecology* 11, 205-221.
- Cause, M. L., Rudder, E. J. and Kynaston, W. T. (1989) *Queensland Timbers: Their Nomenclature, Density and Lyctid-susceptibility*. Technical pamphlet no. 2, Queensland Department of Forestry, Queensland, Indooroopilly.
- Chazdon, R. L. (2003) Tropical forest recovery: legacies of human impact and natural disturbances. *Perspectives in Plant Ecology, Evolution and Systematics* 6, 51-71.
- Chen, J., Franklin, J. F. and Spies, T. A. (1992) Vegetation responses to edge environments in old-growth Douglas-fir forests. *Ecological Applications* 2, 387-396.
- Chen, J., Franklin, J. F. and Spies, T. A. (1995) Growing-season microclimatic gradients from clearcut edges into old-growth Douglas-fir forests. *Ecological Applications* 5, 74-86.
- Cochrane, M. A. and Laurance, W. F. (2002) Fire as a large-scale edge effect in Amazonian forests. *Journal of Tropical Ecology* 18, 311-325.
- Cooper, W. and Cooper, W. T. (1994) *Fruits of the Rain Forest: A Guide to Fruits in Australian tropical Rain Forests*. GEO Productions, Chatswood, NSW.
- D'Angelo, S. A., Andrade, A. C. S., Laurance, S. G., Laurance, W. F. and Mesquita, R. C. G. (2004) Inferred causes of tree mortality in fragmented and intact Amazonian forests. *Journal of Tropical Ecology* 20, 243-246.

- Essen, P. -A. (1994) Tree mortality patterns after experimental fragmentation of an old-growth conifer forest. *Biological Conservation* 68, 19-28.
- Everham, E.M. and Brokaw, N.V.L. (1996) Forest Damage and Recovery from Catastrophic Wind. *The Botanical Review* 62: 113-185.
- Ferreira, L. V. and Laurance, W. F. (1997) Effects of forest fragmentation on mortality and damage of selected trees in central Amazonia. *Conservation Biology* 11, 797-801.
- Floyd, A. G. (1989) *Rainforest Trees of Mainland South-eastern Australia*. Inkata Press, Melbourne.
- Falster, D. S. (2006) Sapling strength and safety: the importance of wood density in tropical forests. *New Phytologist* 171, 237-239.
- Falster, D. S. and Westoby, M. (2005) Alternative height strategies among 45 dicot rain forest species from tropical Queensland, Australia. *Journal of Ecology* 93, 521-535.
- Franklin, J., Drake, D. R., McConkey, K. R., Tonga, F. and Smith, L. B. (2004) The effects of Cyclone *Waka* on the structure of lowland tropical rain forest in Vava'u, Tonga. *Journal of Tropical Ecology* 20, 409-420.
- Gascon, C., Williamson, G. B. and da Fonseca, G. A. B. (2000) Receding Forest Edges and Vanishing Reserves. *Science* 288, 1356-1358.
- Goosem, M. (2006) Weed surveys along highways, roads and powerline clearings traversing the Wet Tropics World Heritage Area. In: *Weed incursions along roads and powerlines in the Wet Tropics World Heritage Area: the potential of remote sensing as an indicator of weed infestations* (eds. M. Goosem and S. M. Turton) pp 1-25. Cooperative Research Centre for Tropical Rainforest Ecology and Management. Rainforest CRC, Cairns. Available online at http://www.jcu.edu.au/rainforest/publications/weed_incursions.htm
- Goosem, M. and Turton, S. M. (2006) *Weed incursions along roads and powerlines in the Wet Tropics World Heritage Area: the potential of remote sensing as an indicator of weed infestations*. Cooperative Research Centre for Tropical Rainforest Ecology and Management. Rainforest CRC, Cairns. Available online at http://www.jcu.edu.au/rainforest/publications/weed_incursions.htm
- Goosem, M., Searle, L. and Walker, S. (in prep.) Remote sensing of severity of vegetation damage caused by severe Tropical Cyclone *Larry* to rainforest canopies in the forest interior and adjacent to a powerline clearing and highway. Marine and Tropical Sciences Research Facility, Cairns.
- Grove, S. J., Turton, S. M. and Siegenthaler, D. T. (2000) Mosaics of canopy openness induced by tropical cyclones in lowland rain forests with contrasting management histories in northeastern Australia. *Journal of Tropical Ecology* 16, 883-894.
- Harper, K. A., Macdonald, S. E., Burton, P. J., Chen, J., Brososke, K. D., Saunders, S. C., Euskirchen, E. S., Roberts, D., Jaiteh, M. S. and Esseen, P. (2005) Edge Influence on Forest Structure and Composition in Fragmented Landscapes. *Conservation Biology* 19, 768-782.
- Hyland, B. P. M. (1989) A Revision of Lauraceae in Australia (excluding *Cassytha*). *Australian Systematic Botany* 2, 135-367.
- Hyland, B. P. M., Whiffin, T., Christophel, D. C., Gray, B. and Elick, R. W. (2003) *Australian Tropical Rain Forest Plants: Trees, Shrubs and Vines*. CSIRO Publishing, Collingwood.

Jackes, B. R. and Cairns, A. (2001) *Plants of the Tropics; Rainforest to Heath; An Identification Guide*. James Cook University, Townsville.

Jones, H. G. (1992) *Plants and microclimate: a quantitative approach to environmental plant physiology*. Second Edition. Cambridge University Press, Cambridge.

Kapos, V., Wandelli, E., Camargo, J. L. and Ganade, G. (1997) Edge-Related Changes in Environment and Plant Resources Due to Forest Fragmentation in Central Amazonia. In: *Tropical Forest Remnants, Ecology, Management, and Conservation of Fragmented Communities* (eds. W. F. Laurance and R. O. Bierregaard) pp. 33-44. The University of Chicago Press, Chicago.

Laurance, W. F. (1991) Edge Effects in Tropical Forest Fragments: Applications of a Model for the Design of Nature Reserves. *Biological Conservation* 57, 205-219.

Laurance, W. F. (1997) Hyper-Disturbed Parks: Edge Effects and the Ecology of Isolated Rainforest Reserves in Tropical Australia. In: *Tropical Forest Remnants, Ecology, Management, and Conservation of Fragmented Communities* (eds. W. F. Laurance and R. O. Bierregaard) pp. 71-82. The University of Chicago Press, Chicago.

Laurance, W. F., Laurance, S. G., Ferreira, L. V., Rankin-de Merona, J. M., Gascon, C. and Lovejoy, T. E. (1997) Biomass Collapse in Amazonian forest fragments. *Science* 278, 1117-1118.

Laurance, W. F., Ferreira, L. V., Rankin-de Merona, J. M. and Laurance, S. G. (1998) Rain forest fragmentation and the dynamics of Amazonian tree communities. *Ecology* 79, 2032-2040.

Laurance, W. F., Delamonica, P., Laurance, S. G., Vasconcelos, H. L. and Lovejoy, T. E. (2000) Rainforest fragmentation kills big trees. *Nature* 404, 836.

Laurance, W. F., Williamson, G. B., Delamonica, P., Oliveira, A., Lovejoy, T. E., Gascon, C. and Pohl, L. (2001) Effects of a strong drought on Amazonian forest fragments and edges. *Journal of Tropical Ecology* 17, 771-785.

Laurance, W. F., Lovejoy, T. E., Vasconcelos, H. L., Bruna, E. M., Didham, R. K., Stouffer, P. C., Gascon, C., Bierregaard, R. O., Laurance, S. G. and Sampaio, E. (2002) Ecosystem Decay of Amazonian Forest Fragments: A 22-Year Investigation. *Conservation Biology* 16, 605-618.

Leigh, E. G., Wright, S. J., Herre, E. A. and Putz, F. E. (1993) The decline of tree diversity on newly isolated tropical islands: a test of a null hypothesis and some implications. *Evolutionary Ecology* 7, 76-102.

Lin, Y., Hulting, M. L. and Augspurger, C. K. (2004) Causes of spatial patterns of dead trees in forest fragments in Illinois. *Plant Ecology* 170, 15-27.

McNaughton, K. G. (1989) Micrometeorology of shelter belts and forest edges. *Philosophical Transactions of the Royal Society of London Series B Biological Sciences* 324, 351-368.

Newmark, W. D. (2001) Tanzanian Forest Edge Microclimatic Gradients: Dynamic Patterns. *Biotropica* 33, 2-11.

Ostertag, R., Silver, W. L. and Lugo, A. E. (2005) Factors Affecting Mortality and Resistance to Damage Following Hurricanes in a Rehabilitated Subtropical Moist Forest. *Biotropica* 37, 16-24.

- Osunkoya, O. O. (1996) Light requirements for regeneration in tropical forest plants: Taxon-level and ecological attribute effects. *Australian Journal of Ecology* 21, 429-441.
- Osunkoya, O. O., Sheng, T. K., Mahmud, N. and Damit, N. (2007) Variation in wood density, wood water content, stem growth and mortality among twenty-seven tree species in a tropical rainforest on Borneo Island. *Austral Ecology* 32, 191-201.
- Pohlman, C. L. (2006) *Internal fragmentation in the rainforest: edge effects of highways, powerlines and watercourses on tropical rainforest understorey microclimate, vegetation structure and composition, physical disturbance and seedling regeneration*. PhD thesis, James Cook University, Cairns.
- Pohlman, C. L., Turton, S. M. and Goosem, M. (2007) Edge Effects of Linear Canopy Openings on Tropical Rainforest Understorey Microclimate. *Biotropica* 39, 62-71.
- Poorter, L., Bongers, L. and Bongers, F. (2006) Architecture of 54 moist-forest tree species: traits, trade-offs, and functional groups. *Ecology* 87, 1289-1301.
- Ray, D., Nepstad, D. and Moutinho, P. (2005) Micrometeorological and canopy controls of fire susceptibility in a forested Amazon landscape. *Ecological Applications* 15, 1664-1678.
- Reynolds, B. E. (1994) *A study of weed (Lantana camara) infestation of tracks through rainforest on the Atherton Tableland, North Queensland*. Honours Thesis, James Cook University.
- Ryall, K. L. and Smith, S. M. (2005) Patterns of damage and mortality in red pine plantations following a major ice storm. *Canadian Journal of Forest Research* 35, 487-493.
- Siegenthaler, S. and Turton, S. M. (2000) Edge effects of roads and powerlines on microclimate. In: *Impacts of Roads and Powerlines on the Wet Tropics of Queensland World Heritage Area*. (eds. M. Goosem, and S. M. Turton) pp. 19-43. Wet Tropics Management Authority and Cooperative Research Centre for Tropical Rainforest Ecology and Management. Available online at http://www.jcu.edu.au/rainforest/publications/impacts_roadspowerlines.htm
- Smith, W. J. ., Kynaston, W. T., Cause, M. L. and Grimmett, J. G. (1991) *Building Timbers: Properties and Recommendations for their Use in Queensland*. Technical Pamphlet No. 1. Queensland Forest Service, Department of Primary Industries, Indooroopilly.
- Tracey, J. G. (1982) *The Vegetation of the Humid Tropical Region of North Queensland*. CSIRO, Melbourne.
- Tucker, N. I. J. (2001) *Vegetation recruitment in a restored habitat linkage in tropical North Queensland*. Masters Thesis, James Cook University, Cairns.
- Tucker, N. I. J. and Murphy, T. M. (1997) The effects of ecological rehabilitation on vegetation recruitment: some observations from the Wet Tropics of North Queensland. *Forest Ecology and Management* 99, 133-152.
- Turner, I. M. (2001) *The Ecology of Trees in the Tropical Rain Forest*. Cambridge University Press, Cambridge, United Kingdom.
- Turton, S. M. (1992) Understorey light environments in a north-east Australian rain forest before and after a tropical cyclone. *Journal of Tropical Ecology* 8, 241-252.

Turton, S. (in prep) Initial effects of Cyclone *Larry* on the forest landscapes of northeast Australia, including comparisons with previous cyclones impacting the region between 1858 and 2006. *Austral Ecology*.

Turton, S. and Freiburger, H. J. (1997) Edge and Aspect Effects on the Microclimate of a Small Tropical Forest Remnant on the Atherton Tableland, Northeastern Australia. In: *Tropical Forest Remnants, Ecology, Management, and Conservation of Fragmented Communities* (eds. W. F. Laurance and R. O. Bierregaard) pp. 45-54. The University of Chicago Press, Chicago.

Turton, S. M. and Siegenthaler, D. T. (2004) Immediate impacts of a severe tropical cyclone on the microclimate of a rain-forest canopy in north-eastern Australia. *Journal of Tropical Ecology* 20, 583-586.

Unwin, G. L., Applegate, G. B., Stocker, G. C. and Nicholson, D. I. (1988) Initial effects of Tropical Cyclone *Winifred* on forests in north Queensland. *Proceedings of the Ecological Society of Australia* 15, 283-296.

van Gelder, H. A., Poorter, L. and Sterck, F. J. (2006) Wood mechanics, allometry, and life-history variation in a tropical rain forest tree community. *New Phytologist* 171, 367-378.

Watson, C. J. J. (1951) *North Queensland Building Timbers and Specifications for Their Use*. Queensland Forest Service Pamphlet No. 1. Government Printer, Brisbane.

Webb, L. J. (1958) Cyclones as an ecological factor in tropical lowland rain-forest, north Queensland. *Australian Journal of Botany* 6, 220-230.

Werren, G. (2001) *Environmental Weeds of the Wet Tropics Bioregion: Risk Assessment and Priority Ranking*. Cooperative Research Centre for Tropical Rainforest Ecology and Management. Rainforest CRC, Cairns. Available at http://www.jcu.edu.au/rainforest/publications/enviro_n_weeds.htm

White, E., Tucker, N., Meyers, N. and Wilson, J. (2004) Seed dispersal to revegetated isolated rainforest patches in North Queensland. *Forest Ecology and Management* 192, 409-426.

Williams, K. A. W. (1984) *Native Plants of Queensland, Volume 2*. National Library of Australia, Canberra.

Williams, K. A. W. (1987) *Native Plants of Queensland, Volume 3*. National Library of Australia, Canberra.

Williams-Linera, G. (1990) Vegetation structure and environmental conditions of forest edges in Panama. *Journal of Ecology* 78, 356-373.

Zimmerman, J. K., Everham, E. M., Waide, R. B., Lodge, D. J., Taylor, C. M. and Brokaw, N. V. L. (1994) Responses of tree species to hurricane winds in subtropical wet forest in Puerto Rico: implications for tropical tree life histories. *Journal of Ecology* 82, 911-922.

Appendix 1

Wood densities of tree species encountered in the vegetation survey

Data obtained from Osunkoya (1996), Cause *et al.* (1989), Hyland (1989) and from expert advice (Dr Steve Goosem, *pers. comm.*, citing Floyd 1989 and Watson 1951).

Species	Family	Wood Density (kg m ⁻³)
<i>Aceratium megalospermum</i>	Eleocharaceae	625
<i>Acmena graveolens</i>	Myrtaceae	595
<i>Acronychia vestita</i>	Rutaceae	705
<i>Aglaia meridionalis</i>	Meliaceae	700 (47) [†]
<i>Aglaia tomentosa</i>	Meliaceae	700 (47) [†]
<i>Alangium villosum</i>	Alangiaceae	705
<i>Alphitonia petriei</i>	Rhamnaceae	515
<i>Alstonia scholaris</i>	Apocynaceae	400
<i>Antirrhoea tenuiflora</i>	Rubiaceae	805 (58) [†]
<i>Apodytes brachystylis</i>	Icacinaceae	655
<i>Archidendron whitei</i>	Mimosaceae	705 (74) ^{†‡}
<i>Argyrodendron peralatum</i>	Sterculiaceae	800
<i>Argyrodendron trifoliolatum</i>	Sterculiaceae	925
<i>Austromyrtus bidwillii</i>	Myrtaceae	775 (21) ^{†§}
<i>Austromyrtus dallachiana</i>	Myrtaceae	775 (21) ^{†§}
<i>Austromyrtus shepherdii</i>	Myrtaceae	775 (21) ^{†§}
<i>Beilschmedia bancroftii</i>	Lauraceae	640
<i>Beilschmedia recurva</i>	Lauraceae	620
<i>Beilschmedia tooram</i>	Lauraceae	850
<i>Beilschmedia volckii</i>	Lauraceae	545
<i>Bischofia javanica</i>	Euphorbiaceae	655
<i>Breynia stipitata</i>	Euphorbiaceae	690 (37) [†]
<i>Brombya platynema</i>	Rutaceae	710
<i>Cananga odorata</i>	Annonaceae	465
<i>Cardwellia sublimis</i>	Proteaceae	560
<i>Carnarvonium araliifolia</i>	Proteaceae	690
<i>Castanospermum australe</i>	Fabaceae	755
<i>Castanospora alphandii</i>	Sapindaceae	705
<i>Celtis paniculata</i>	Ulmaceae	705
<i>Chionanthus axillaris</i>	Oleaceae	935 (40) [†]
<i>Chisocheton longistipitatus</i>	Meliaceae	545

Species	Family	Wood Density (kg m ⁻³)
<i>Cinnamomum laubatii</i>	Lauraceae	480
<i>Citronella smythii</i>	Icacinaceae	675
<i>Clerodendron grayi</i>	Verbenaceae	585 (37) [†]
<i>Corynocarpus cribbianus</i>	Corynocarpaceae	690
<i>Cryptocarya angulata</i>	Lauraceae	755
<i>Cryptocarya corrugata</i>	Lauraceae	800
<i>Cryptocarya grandis</i>	Lauraceae	830
<i>Cryptocarya mackinnoniana</i>	Lauraceae	880
<i>Cryptocarya melanocarpa</i>	Lauraceae	775
<i>Cryptocarya murrayi</i>	Lauraceae	785
<i>Cryptocarya oblata</i>	Lauraceae	560
<i>Cryptocarya pleurosperma</i>	Lauraceae	690
<i>Daphnandra repandula</i>	Monimiaceae	675
<i>Davidsonia pruriens</i>	Davidsoniaceae	875
<i>Diospiros cupulosa</i>	Ebenaceae	1010 (122) [†]
<i>Diospiros</i> sp. "twice as flat"	Ebenaceae	1010 (122) [†]
<i>Diploglottis bracteata</i>	Sapindaceae	995
<i>Diploglottis smithii</i>	Sapindaceae	830 (22) [†]
<i>Doryphora aromatica</i>	Monimiaceae	560
<i>Dysoxylum klanderi</i>	Meliaceae	945
<i>Dysoxylum oppositifolium</i>	Meliaceae	880
<i>Dysoxylum papuanum</i>	Meliaceae	735
<i>Dysoxylum pettigrewianum</i>	Meliaceae	865
<i>Elaeocarpus grandis</i>	Elaeocarpaceae	495
<i>Elaeocarpus largiflorens</i>	Elaeocarpaceae	450
<i>Endiandra bessaphila</i>	Lauraceae	665
<i>Endiandra compressa</i>	Lauraceae	995
<i>Endiandra globosa</i>	Lauraceae	915
<i>Endiandra insignis</i>	Lauraceae	750
<i>Endiandra leptodendron</i>	Lauraceae	870
<i>Endiandra monothyra</i>	Lauraceae	800
<i>Endiandra palmerstonii</i>	Lauraceae	690
<i>Endiandra sankeyana</i>	Lauraceae	755
<i>Endiandra sideroxylon</i>	Lauraceae	800
<i>Ficus congesta</i>	Moraceae	350
<i>Ficus copiosa</i>	Moraceae	350
<i>Ficus crassipes</i>	Moraceae	350
<i>Ficus leptoclada</i>	Moraceae	560

Species	Family	Wood Density (kg m ⁻³)
<i>Ficus pleurocarpa</i>	Moraceae	470
<i>Ficus septica</i>	Moraceae	350
<i>Ficus variegata</i>	Moraceae	400
<i>Ficus virens</i> var. <i>virens</i>	Moraceae	400
<i>Flindersia acuminata</i>	Rutaceae	530
<i>Flindersia brayleyana</i>	Rutaceae	575
<i>Franciscodendron laurifolium</i>	Sterculiaceae	450
<i>Gardenia ovularis</i>	Rubiaceae	850
<i>Gessios biagiana</i>	Cunoniaceae	640
<i>Gillbeea adenopetala</i>	Cunoniaceae	530
<i>Glochidion harveyanum</i>	Euphorbiaceae	785
<i>Glochidion sumatrum</i>	Euphorbiaceae	705
<i>Guioa lasioneura</i>	Sapindaceae	830 (22) [†]
<i>Haplostichanthus</i> sp. Johnstone River LWJ 471	Annonaceae	565 (38) [†]
<i>Helicia nortoniana</i>	Proteaceae	725 (33) ^{†¶}
<i>Hollandaea sayeriana</i>	Proteaceae	725 (33) ^{†¶}
<i>Hylandia dockrillii</i>	Euphorbiaceae	560
<i>Irvingbaileya australis</i>	Icacinaceae	495
<i>Levieria acuminata</i>	Monimiaceae	435
<i>Litsea leefeana</i>	Lauraceae	480
<i>Macaranga inamoena</i>	Euphorbiaceae	560
<i>Mallotus paniculatus</i>	Euphorbiaceae	690 (37) [†]
<i>Melicope bonwickii</i>	Rutaceae	465
<i>Melicope elleryana</i>	Rutaceae	610
<i>Melicope vitiflora</i>	Rutaceae	625
<i>Melicope xanthoxyloides</i>	Rutaceae	495
<i>Mischocarpus lachnocarpus</i>	Sapindaceae	830 (22) [†]
<i>Myristica insipida</i>	Myristicaceae	560
<i>Neolitsea dealbata</i>	Lauraceae	680
<i>Niemeyera prunifera</i>	Sapotaceae	610
<i>Omalanthus novo-guineensis</i>	Euphorbiaceae	320
<i>Opistheolepis heterophylla</i>	Proteaceae	610
<i>Ostrearia australiana</i>	Hamamelidaceae	755
<i>Palaquium galatoxylon</i>	Sapotaceae	560
<i>Phaleria clerodendron</i>	Thymelaeaceae	655 (16) [‡]
<i>Pilidiostigma tropicum</i>	Myrtaceae	775 (21) [†]
<i>Pitiviaster haplophylla</i>	Rutaceae	835

Species	Family	Wood Density (kg m ⁻³)
<i>Podocarpus dispersus</i>	Podocarpaceae	580 (45) [†]
<i>Polyalthia michaelii</i>	Annonaceae	625
<i>Polyosma hirsute</i>	Grossulariaceae	720 (na) [†]
<i>Polyscias australiana</i>	Araliaceae	575
<i>Polyscias elegans</i>	Araliaceae	480
<i>Polyscias murrayi</i>	Araliaceae	400
<i>Pouteria castanosperma</i>	Sapotaceae	975
<i>Prunus turneriana</i>	Rosaceae	530
<i>Pseuduvaria villosa</i>	Annonaceae	565 (38) [†]
<i>Rhodamnia sessiliflora</i>	Myrtaceae	975
<i>Rhodomyrtus pervigata</i>	Myrtaceae	775 (21) [†]
<i>Rhysotoechia robertsonii</i>	Sapindaceae	830 (22) [†]
<i>Rockinghamia angustifolia</i>	Euphorbiaceae	800
<i>Sarcotoechia protracta</i>	Sapindaceae	830 (22) [†]
<i>Schefflera actinophylla</i>	Araliaceae	480
<i>Siphonodon membranaceus</i>	Celastraceae	835
<i>Sloanea australis</i>	Eleocarpaceae	625
<i>Sloanea macbrydei</i>	Eleocarpaceae	575
<i>Symplocus cochinchinensis</i>	Symplocaceae	545
<i>Symplocus paucistaminea</i>	Symplocaceae	585 (40) [†]
<i>Synima cordierorum</i>	Sapindaceae	945
<i>Synima macrophylla</i>	Sapindaceae	830 (22) [†]
<i>Synuom glandulosum</i> ssp. <i>paniculosum</i>	Meliaceae	675
<i>Synuom muelleri</i>	Meliaceae	625
<i>Syzygium alliligineum</i>	Myrtaceae	610
<i>Syzygium cormiflorum</i>	Myrtaceae	770
<i>Syzygium gustavioides</i>	Myrtaceae	690
<i>Syzygium sayeri</i>	Myrtaceae	840
<i>Tetrasynandra laxiflora</i>	Monimiaceae	640
<i>Toechima erythrocarpum</i>	Sapindaceae	785
<i>Toechima monticola</i>	Sapindaceae	830 (22) [†]
<i>Xanthophyllum octandrum</i>	Xanthophyllaceae	800

[†] Family average used (obtained from Cause *et al.* 1989 and Smith *et al.* 1991). Standard deviation is shown in parentheses.

[‡] Mimosaceae average excludes the genus *Acacia*.

[§] Myrtaceae average excludes the genera *Corymbia*, *Eucalyptus*, *Leptospermum*, *Lophostemon* and *Melaleuca*.

[¶] Proteaceae average excludes the genera *Banksia* and *Grevillea*.

[¿] No data available on the family Thymelaeaceae; dataset mean used instead.

Further information

Marine and Tropical Sciences Research Facility
PO Box 1762
CAIRNS QLD 4870

or

Marine and Tropical Sciences Research Facility
PO Box 772
TOWNSVILLE QLD 4810

This document is available for download at <http://www.rrrc.org.au/publications>

Credits: **Southern cassowary** Wet Tropics Management Authority; **Hill Inlet in the Whitsundays** Department of Foreign Affairs and Trade - Overseas Information Branch; **Butterfly fish** Robert Thorn; **Rainforest fruits** Wet Tropics Management Authority; **Cover image strip - fieldwork and cyclone damage** Catherine Pohlman.